

FORM PTO-1390 (Modified) (REV 11-2000)		U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE		ATTORNEY'S DOCKET NUMBER
TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. 371				29342/36539A
INTERNATIONAL APPLICATION NO. PCT/US00/20981		INTERNATIONAL FILING DATE 01 August 2000		U.S. APPLICATION NO. (IF KNOWN, SEE 37 CFR 10/031463
				PRIORITY DATE CLAIMED 03 August 1999
TITLE OF INVENTION BETA-CARBOLINE DRUG PRODUCTS				
APPLICANT(S) FOR DO/EO/US ANDERSON, Neil R.; HARTAUER, Kerry J.; KRAL, Martha A.; STEPHENSON, Gregory A.				
Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:				
<ol style="list-style-type: none"> <input checked="" type="checkbox"/> This is a FIRST submission of items concerning a filing under 35 U.S.C. 371. <input type="checkbox"/> This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. 371. <input type="checkbox"/> This is an express request to begin national examination procedures (35 U.S.C. 371(f)). The submission must include items (5), (6), (9) and (24) indicated below. <input checked="" type="checkbox"/> The US has been elected by the expiration of 19 months from the priority date (Article 31). <input checked="" type="checkbox"/> A copy of the International Application as filed (35 U.S.C. 371 (c) (2)) <ol style="list-style-type: none"> <input type="checkbox"/> is attached hereto (required only if not communicated by the International Bureau). <input type="checkbox"/> has been communicated by the International Bureau. <input checked="" type="checkbox"/> is not required, as the application was filed in the United States Receiving Office (RO/US). <input type="checkbox"/> An English language translation of the International Application as filed (35 U.S.C. 371(c)(2)). <ol style="list-style-type: none"> <input type="checkbox"/> is attached hereto. <input type="checkbox"/> has been previously submitted under 35 U.S.C. 154(d)(4). <input checked="" type="checkbox"/> Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371 (c)(3)) <ol style="list-style-type: none"> <input type="checkbox"/> are attached hereto (required only if not communicated by the International Bureau). <input type="checkbox"/> have been communicated by the International Bureau. <input type="checkbox"/> have not been made; however, the time limit for making such amendments has NOT expired. <input checked="" type="checkbox"/> have not been made and will not be made. <input type="checkbox"/> An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)). <input type="checkbox"/> An oath or declaration of the inventor(s) (35 U.S.C. 371 (c)(4)). <input type="checkbox"/> An English language translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371 (c)(5)). <input checked="" type="checkbox"/> A copy of the International Preliminary Examination Report (PCT/IPEA/409). <input checked="" type="checkbox"/> A copy of the International Search Report (PCT/ISA/210). 				
Items 13 to 20 below concern document(s) or information included:				
<ol style="list-style-type: none"> <input type="checkbox"/> An Information Disclosure Statement under 37 CFR 1.97 and 1.98. <input type="checkbox"/> An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included. <input checked="" type="checkbox"/> A FIRST preliminary amendment. <input type="checkbox"/> A SECOND or SUBSEQUENT preliminary amendment. <input type="checkbox"/> A substitute specification. <input type="checkbox"/> A change of power of attorney and/or address letter. <input type="checkbox"/> A computer-readable form of the sequence listing in accordance with PCT Rule 13ter.2 and 35 U.S.C. 1.821 - 1.825. <input type="checkbox"/> A second copy of the published international application under 35 U.S.C. 154(d)(4). <input type="checkbox"/> A second copy of the English language translation of the international application under 35 U.S.C. 154(d)(4). <input checked="" type="checkbox"/> Certificate of Mailing by Express Mail <input checked="" type="checkbox"/> Other items or information: Return receipt postcard 				

U.S. APPLICANT 10/031463	INTERNATIONAL APPLICATION NO. PCT/US00/20981	ATTORNEY'S DOCKET NUMBER 29342/36539A			
24. The following fees are submitted:		CALCULATIONS PTO USE ONLY			
BASIC NATIONAL FEE (37 CFR 1.492 (a) (1) - (5)) :					
<input type="checkbox"/> Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO \$1040.00 <input checked="" type="checkbox"/> International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO \$890.00 <input type="checkbox"/> International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.445(a)(2)) paid to USPTO \$740.00 <input type="checkbox"/> International preliminary examination fee (37 CFR 1.482) paid to USPTO but all claims did not satisfy provisions of PCT Article 33(1)-(4) \$710.00 <input type="checkbox"/> International preliminary examination fee (37 CFR 1.482) paid to USPTO and all claims satisfied provisions of PCT Article 33(1)-(4) \$100.00					
ENTER APPROPRIATE BASIC FEE AMOUNT =		\$890.00			
Surcharge of \$130.00 for furnishing the oath or declaration later than months from the earliest claimed priority date (37 CFR 1.492 (c)).		<input type="checkbox"/> 20 <input type="checkbox"/> 30 \$0.00			
CLAIMS		NUMBER FILED	NUMBER EXTRA	RATE	
Total claims		19 - 20 =	0	x \$18.00	\$0.00
Independent claims		7 - 3 =	4	x \$84.00	\$336.00
Multiple Dependent Claims (check if applicable).				<input type="checkbox"/>	\$0.00
TOTAL OF ABOVE CALCULATIONS =		\$1,226.00			
<input type="checkbox"/> Applicant claims small entity status. See 37 CFR 1.27). The fees indicated above are reduced by 1/2.		\$0.00			
SUBTOTAL =		\$1,226.00			
Processing fee of \$130.00 for furnishing the English translation later than months from the earliest claimed priority date (37 CFR 1.492 (f)).		<input type="checkbox"/> 20 <input type="checkbox"/> 30	+ \$0.00		
TOTAL NATIONAL FEE =		\$1,226.00			
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31) (check if applicable).				<input type="checkbox"/>	\$0.00
TOTAL FEES ENCLOSED =		\$1,226.00			
				Amount to be: refunded	\$
				charged	\$
a. <input checked="" type="checkbox"/> A check in the amount of \$1,226.00 to cover the above fees is enclosed. b. <input type="checkbox"/> Please charge my Deposit Account No. _____ in the amount of _____ to cover the above fees. A duplicate copy of this sheet is enclosed. c. <input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 13-2855 A duplicate copy of this sheet is enclosed. d. <input type="checkbox"/> Fees are to be charged to a credit card. WARNING: Information on this form may become public. Credit card information should not be included on this form. Provide credit card information and authorization on PTO-2038.					
NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.					
SEND ALL CORRESPONDENCE TO: NAPOLI, James J. Customer No. 04743 Marshall, Gerstein & Borun 6300 Sears Tower 233 South Wacker Drive Chicago, Illinois 60606 United States of America					
 SIGNATURE James J. Napoli NAME 32,361 REGISTRATION NUMBER 17 January 2002 DATE					

3 00031463 107031463
JC13 Rec'd PCT/PTO 17 JAN 2002

PATENT APPLICATION

IN THE UNITED STATES PATENT
AND TRADEMARK OFFICE

Richard J.
Richard Zimmermann

**PRELIMINARY AMENDMENT ACCOMPANYING
NEW APPLICATION TRANSMITTAL**

Box PCT
Commissioner of Patents
Washington, D.C. 20231

Sir:

Please amend the above-identified application filed under 35 U.S.C. §371 as follows:

IN THE SPECIFICATION:

Page 1, after the title, please delete the CROSS-REFERENCE TO RELATED APPLICATION in its entirety and insert therefor:

--CROSS-REFERENCE TO RELATED APPLICATIONS


This is the U.S. national phase application of International Application No. PCT/US00/20981, filed on August 1, 2000, which claims the benefit of provisional patent application Serial No. 60/147,048, filed August 3, 1999.--

IN THE CLAIMS:

Cancel claims 17-19, without prejudice.

Add new claims 20-22 as follows:

--20. A method of treating sexual dysfunction in a patient in need thereof, which comprises administering to the patient a therapeutically effective amount of a free form of a compound having a formula



which exhibits a C_{max} of about 180 to about 280 micrograms/liter and an AUC (0-24) of about 2280 to about 3560 micrograms hour/liter, measured using a 10 milligram dose of the compound.

21. The method of claim 20 wherein the sexual dysfunction is male erectile dysfunction.

22. The method of claim 20 wherein the sexual dysfunction is female sexual arousal disorder.--

REMARKS

Claims 1-19 are pending in the application. Claims 17-19 have been cancelled by this amendment. Claims 20-22 have been added. Therefore, claims 1-16 and 20-22 are at issue.

The amendments are described in more detail below. Pursuant to 37 C.F.R. §1.121, a marked-up version of the changes made to the specification and claims by the present amendment is attached hereto following the signature page of this amendment. The first page of the marked-up version of the changes is captioned "Version With Markings to Show Changes Made."

This preliminary amendment adds no new matter. The specification has been amended to insert a new cross reference to related applications. The claims have been amended to conform the claims to U.S. practice. Support for new claims 20-22 can be found in originally filed claims 7-9 and 18.


It is submitted that this amendment should be entered and that the claims are in proper form for examination. An early and favorable action on the merits is respectfully requested.

Should the examiner wish to discuss the foregoing or any matter of form in an effort to advance this application toward allowance, the examiner is urged to telephone the undersigned at the indicated number.

Respectfully submitted,

MARSHALL, GERSTEIN & BORUN

By



James J. Napoli
(Registration No. 32,361)
Attorneys for Applicants
6300 Sears Tower
233 South Wacker Drive
Chicago, Illinois 60606
(312) 474-6300

Chicago, Illinois
January 17, 2002

VERSION WITH MARKINGS TO SHOW CHANGES MADE
(U.S. National Stage of PCT/US00/11130
filed January 17, 2002)

IN THE SPECIFICATION:

The following cross-reference to related application has been inserted into the specification:

CROSS-REFERENCE TO RELATED APPLICATIONS


This is the U.S. national phase application of International Application No. PCT/US00/20981, filed on April 26, 2000, which claims the benefit of provisional patent application Serial No. 60/147,048, filed August 3, 1999.

IN THE CLAIMS:

Claims 17-19 have been cancelled without prejudice.

New claims 20-22 have been added to the application.

20. A method of treating sexual dysfunction in a patient in need thereof, which comprises administering to the patient a therapeutically effective amount of a free form of a compound having a formula



which exhibits a C_{max} of about 180 to about 280 micrograms/liter and an AUC (0-24) of about 2280 to about 3560 micrograms hour/liter, measured using a 10 milligram dose of the compound.

21. The method of claim 20 wherein the sexual dysfunction is male erectile dysfunction.

22. The method of claim 20 wherein the sexual dysfunction is female sexual arousal disorder.

β-CARBOLINE DRUG PRODUCTS

CROSS REFERENCE TO RELATED APPLICATIONS

5 This application claims the benefit of
provisional patent application Serial No.
60/147,048, filed August 3, 1999.

FIELD OF THE INVENTION

10 The present invention relates to the
fields of pharmaceutical and organic chemistry, and
to a β-carboline compound which is useful for the
treatment of various medical indications where
15 inhibition of type 5 cGMP-specific phosphodiesterase
(PDE5) is desired. More particularly the present
invention provides a free drug form of β-carboline
particles in a size range allowing for uniform
formulation of stable pharmaceutical compositions,
20 especially compositions providing desired bioavail-
ability properties heretofore not provided in the
art.

BACKGROUND OF THE INVENTION

25 The biochemical, physiological, and clin-
ical effects of cyclic guanosine 3',5'-monophosphate
specific phosphodiesterase (cGMP-specific PDE) in-
hibitors suggest their utility in a variety of
30 disease states in which modulation of smooth muscle,
renal, hemostatic, inflammatory, and/or endocrine
function is desired. Type 5 cGMP-specific phospho-
diesterase (PDE5) is the major cGMP hydrolyzing

- 2 -

enzyme in vascular smooth muscle, and its expression in penile corpus cavernosum has been reported (Taher et al., *J. Urol.*, 149:285A (1993)). Thus, PDE5 is an attractive target in the treatment of sexual dysfunction (Murray, *DN&P* 6(3):150-156 (1993)).

Daugan U.S. Patent No. 5,859,006 discloses a class of β -carboline compounds, and pharmaceutical compositions containing the β -carbolines, which are useful in the treatment of conditions wherein inhibition of PDE5 is desired. PCT publication 10 WO 97/03675 discloses use of this class of β -carboline compounds in the treatment of sexual dysfunction.

The poor solubility of many β -carboline compounds useful as PDE5 inhibitors prompted the development of coprecipitate preparations, as disclosed in PCT publication WO 96/38131 and Butler U.S. Patent No. 5,985,326. Briefly, coprecipitates of a β -carboline with polymeric hydroxypropylmethyl-cellulose phthalate, for example, were prepared, milled, mixed with excipients, and compressed into tablets for oral administration. Studies revealed, however, that difficulties arose in generating precisely reproducible lots of coprecipitate product, which makes use of coprecipitates less than ideal in pharmaceutical formulations.

Additionally, clinical studies involving administration of coprecipitate tablets preliminarily revealed that maximum blood concentration of the β -carboline compound is achieved in 3 to 4 hours, with the average time for onset of therapeutic effect not yet precisely determined. In the treatment of sexual dysfunction, such as male erectile

- 3 -

dysfunction or female sexual arousal disorder, however, a more rapid achievement of maximum blood concentration, along with a greater prospect for rapid onset of therapeutic effect, frequently is sought by individuals desiring more immediate and/or less prolonged effects. Accordingly, a need in the art continues to exist for orally administrable β -carboline compounds and β -carboline-containing pharmaceutical compositions having an ability to provide a therapeutic effect within a desirable, or at least acceptable, time frame.


SUMMARY OF THE INVENTION

The present invention provides particulate preparations of a free drug form of a β -carboline compound having specific and defined particle size characteristics. The defined particle size permits a uniform formulation of stable pharmaceutical compositions. In particular, the present invention provides compositions that exhibit a rapid achievement of maximum blood concentration of PDE5 inhibitor and/or a rapid onset of a therapeutic PDE5 inhibitory effect.

The present invention provides a compound having the formula (I)

- 4 -

5



10

(I)

and pharmaceutically acceptable salts and solvates thereof, wherein the compound is a free drug in particulate form, and wherein at least 90% of the particles have a particle size of less than about 40 microns, and preferably less than 30 microns. Highly preferred particulate forms of the β -carboline compound (I) have at least 90% of the particles less than 25 microns in size. Most preferred forms of the free compound (I) are those wherein 90% of the particles are less than 10 microns in size.

The present invention provides, therefore, a free form of a β -carboline compound, and compositions containing the β -carboline compound, which can be used in an effective therapy for conditions wherein inhibition of PDE5 provides a benefit. The free form of β -carboline compound (I) has a particle size such that the onset of beneficial effects of PDE5 inhibition are exhibited in a relatively short time after oral administration.

The present invention further relates to pharmaceutical compositions comprising the particulate compound (I) and one or more pharmaceutically

- 5 -

acceptable carriers, diluents, or excipients. The invention further provides the use of compound (I) and pharmaceutical compositions for treatment of sexual dysfunction, e.g., male erectile dysfunction and female sexual arousal disorder.

Alternatively stated, the present invention provides for the use of the above-described particulate forms of compound (I) for the manufacture of medicaments for the treatment of sexual dysfunction. Specific conditions that can be treated by the compound and compositions of the present invention include, but are not limited to, male erectile dysfunction and female sexual dysfunction, for example, female arousal disorder, also known as female sexual arousal disorder.

Accordingly, one aspect of the present invention is to provide a free drug particulate form of a compound (I), and pharmaceutically acceptable salts and solvates thereof, comprising particles of the compound wherein at least 90% of the particles have a particle size of less than about 40 microns.

Another aspect of the present invention is to provide a pharmaceutical composition comprising particles of the free drug particulate form of compound (I) having a d90 less than 40, and one or more pharmaceutically-acceptable carriers, diluents, or excipients, and a method of manufacturing the composition.

Yet another aspect of the present invention is to provide a method of treating sexual dysfunction in patients in need thereof comprising administering to a patient in need thereof a therapeutically effective amount of a composition

- 6 -

comprising particles of the free drug particulate form of compound (I) having a d90 less than 40 and one or more pharmaceutically-acceptable carriers, diluents, or excipients. The sexual dysfunction can 5 be male erectile dysfunction or female arousal disorder, for example.

Still another aspect of the present invention is to provide a pharmaceutical composition comprising: (a) a free drug form of compound (I), and pharmaceutically-acceptable salts and solvates thereof, and (b) one or more pharmaceutically-acceptable carriers, diluents, or excipients, wherein the composition exhibits a C_{max} of about 180 to about 280 micrograms/liter or an AUC (0-24) of 10 about 2280 to about 3560 microgram · hour/ liter, measured using a 10 milligram dose of the compound. The composition can be a solid, a suspension, or a 15 solution.

Another aspect of the present invention is 20 to provide a pharmaceutical composition comprising: (a) compound (I) and pharmaceutically-acceptable salts and solvates thereof, and (b) one or more pharmaceutically-acceptable carriers, diluents, or excipients, wherein the composition exhibits a C_{max} of about 180 to about 280 micrograms/liter and an 25 AUC (0-24) of about 2280 to about 3560 micrograms · hour/liter, measured using a 10 milligram dose of the compound. The composition can be a solid or a suspension.

Another aspect of the present invention is 30 to provide a pharmaceutical composition comprising: (a) a free drug form of compound (I), and pharmaceutically acceptable salts and solvates

- 7 -

thereof, wherein at least 90% of the particles have a particle size of less than about 10 microns, and (b) one or more pharmaceutically-acceptable carriers, diluents, or excipients, and bioequivalent compositions thereof. The composition can be a solid or a suspension.

These and other aspects of the present invention will become apparent from the following detailed description of the invention.

10

BRIEF DESCRIPTION OF THE DRAWINGS

15 Fig. 1 contains plots of % of dissolved compound (I) vs. time, and illustrates the *in vitro* dissolution characteristics of compound (I) in a varying particle size.

DETAILED DESCRIPTION OF THE INVENTION

20 For purposes of the claimed invention as disclosed and described herein, the following terms and abbreviations are defined as follows.

25 The term "treatment" includes preventing, lowering, stopping, or reversing the progression or severity of a condition or symptoms being treated. As such, the present invention includes both therapeutic and prophylactic administration, as appropriate.

30 The term "effective amount" is an amount of compound (I), or a composition containing compound (I), that is effective in treating the condition or symptom of interest. An effective amount of a compound (I) to treat sexual dysfunction in a male

- 8 -

is an amount sufficient to provide and sustain an erection capable of penetrating the partner. An effective amount of a compound (I) to treat female sexual dysfunction, particularly female sexual arousal disorder, is an amount sufficient to enhance the ability of a female to achieve or sustain an aroused state.

5 The term "free drug" refers to solid particles of compound (I) not intimately embedded in a 10 polymeric coprecipitate.

10 The term "suspension" refers to a liquid 15 composition containing free drug particles of compound (I). The term "solution" refers to a liquid composition having compound (I) dissolved therein.

The term "solvate" comprises one or more molecules of compound (I) associated one or more molecule of a solvent, e.g., water or acetic acid.

20 The term "oral dosage form" is used in a general sense to refer to pharmaceutical products administered via the mouth. Solid oral dosage forms are recognized by those skilled in the art to include such forms as tablets, capsules, and aerosols.

25 The term "pharmaceutically acceptable" means carriers, excipients, diluents, salt forms of compound (I), and other formulation ingredients that are compatible with all other ingredients of a composition, and are not deleterious to an 30 individual treated with the composition.

The nomenclature describing the particle size of compound (I) is commonly referred to, and is herein, as the "d90." For example, a d90 of 40 (or

- 9 -


d₉₀=40) means that at least 90% of the particles have a particle size of less than 40 microns.

As noted, the present invention provides a compound of structural formula (I), and pharmaceutically acceptable salts and solvates thereof, characterized in that the compound is a free drug in particulate form, wherein at least 90% of the particles have a particle size of less than about 40 microns.

It has been found that by processing (6R-trans)-6-(1,3-benzodioxol-5-yl)-2,3,6,7,12,12a-hexahydro-2-methylpyrazino[1',2':1,6]pyrido[3,4-b]indole-1,4-dione, alternatively named (6R,12aR)-2,3,6,7,12,12a-hexahydro-2-methyl-6-(3,4-methylene-dioxyphenyl)pyrazino[2',1':6.1]pyrido[3,4-b]indole-1,4-dione, as disclosed in Daugan U.S. Patent No. 5,859,006, and represented by structural formula (I):

20

25



(I)

30

- 10 -

5 to bring the particle size within a particular narrow range, manufacturing capability is enhanced, and pharmaceutical compositions can be prepared that exhibit an improved bioavailability of the active ingredient, i.e., compound (I).

10 The present invention encompasses particles of free compound (I) wherein at least 90% of the particles of free drug have a particle size of less than about 40 microns (i.e., $d_{90}=40$), and preferably less than 30 microns. More preferably, at least 90% of the particles have a particle size of less than 25 microns, still more preferably less than 15 microns, and to achieve the full advantage 15 of the present invention, d_{90} is less than 10 microns. Particles having a d_{90} in the nanometer range (e.g., about 200 nm or less, or about 50 nm or less) also are contemplated. However, nanometer sized particles of compound (I) are difficult to 20 handle and to formulate, and tend to aggregate. Therefore, a preferred d_{90} range for the particles of free compound (I) is about 1 to about 40 microns.

25 Preferably, the free drug is crystalline. However, amorphous and partially amorphous forms of compound (I) also are contemplated, and are included within the present invention.

30 It is understood by those familiar with comminution process techniques that the limit set on the size of 90% or more of the particles, using normal milling techniques, is a feature to further distinguish the particulate compounds of the invention from particles exhibiting a broader size

- 11 -

5 distribution. Because of the variation in size encountered in all matter reduced in size by a comminution process, expressing differences in particle size in the manner described herein is readily accepted by those skilled in the art.

10 The present invention also provides pharmaceutical compositions comprising said particulate compound (I) and one or more pharmaceutically acceptable excipients, diluents, or carriers. The excipient, diluent, or carrier can be a solid component of the composition or a liquid component. Accordingly, pharmaceutical compositions containing particles of free compound (I) can be a solid composition, or can be a suspension of free 15 compound (I) particles in a liquid excipient, diluent, or carrier.

20 The compound of the structural formula (I) can be made according to established procedures, such as those detailed in U.S. Patent No. 5,859,006, incorporated herein by reference. The preparation of the compound of structural formula (I) is 25 specifically provided in U.S. Patent No. 5,859,006.

25 Methods of determining the size of particles are well known in the art. For example, the general method of U.S. Patent No. 4,605,517, incorporated herein by reference, could be employed. The following is a description of one nonlimiting method.

30 In preparing the particulate compound of the present invention, a compound of structural formula (I), in its raw state, first is characterized for size using an instrument adapted to measure equivalent spherical volume diameter, e.g., a Horiba

- 12 -

5 LA910 Laser Scattering Particle Size Distribution Analyzer or equivalent instrument. Typically, a representative sample of a compound of structural formula (I) is expected to comprise, in its raw state, particles having a d₉₀ equivalent spherical volume diameter of about 75 to about 200 microns, and with a broad size distribution.

10 After being characterized for size in its raw state, the free drug compound then is milled, for example using a pin mill under suitable conditions of mill rotation rate and feed rate, to bring the particle size value within the above mentioned limits of the present invention. The efficiency of the milling is monitored by sampling, using a Horiba 15 LA910 Laser Scattering Particle Size Distribution Analyzer, and the final particle size is confirmed in a similar manner. If a first pass through the mill fails to produce the required size distribution, then one or more further passes are effected. 20 Other methodologies to prepare particles as described herein are readily available, including a variety of milling techniques, such as hammer or fluid energy mills.

25 The particles of compound (I) in the raw state, as well as after milling or other particle size reduction techniques, are irregular in shape. Therefore, it is necessary to characterize the particles by a measurement different from actual size, like thickness or length, for example, by 30 measurement of a property, like intensity and angle of diffracted light, and equate that measurement to the diameter of known spherical particles having the measured same property. The particles are thus

- 13 -

5 allocated an "equivalent spherical diameter." The values found from characterizing a large number of "unknown" particles can be plotted as cumulative frequency vs. diameter, or in other methods weight vs. diameter, usually adopting percentage undersize values for cumulative frequency or weight. This provides a characteristic curve representing size distribution of the sample, i.e., cumulative percentage undersize distribution curve. Values can be 10 read directly from the curve, or, alternatively, the measurements are plotted on log-probability paper to give a straight line, and the values can be read therefrom.

15 The d₉₀ equivalent spherical volume diameter thus found is a statistical representation of the 90% point on a cumulative frequency plot. As indicated, the d₉₀ equivalent sphere volume diameter of the particles of the milled compound of formula (I) are evaluated using a Horiba LA910 Laser 20 Scattering Particle Size Distribution Analyzer or other such equipment recognized by those skilled in the art. Using such instrument values for a suspension of the particles of unknown size are obtained, and the instrument is monitored using a control 25 sample having particles within the size range expected based on statistical analysis of the control sample.

30 The particle size of compound (I) prior to formulation into a pharmaceutical composition can be measured, for example, as follows. The laser scattering particle size distribution analysis is effected on a small sample of the reduced material, which is suspended in approximately 180 ml of

- 14 -

dispersant solution. Prior to sample suspension, a dispersant solution containing 0.1% SPAN 80 in cyclohexane, and presaturated with compound (I), is prepared. The dispersant solution is filtered 5 through a 0.2 micron microporous membrane filter to provide a particle-free dispersant solution. The sample then is added to the dispersant solution until an acceptable level of laser light obscuration is achieved, at which point the particle size distribution is measured.

10 TriPLICATE measurements are effected as a minimum a) to provide more reliable measurements and b) to check the equivalent sampling of the suspended material. The results are automatically recorded 15 and displayed graphically to give a cumulative % undersize vs. diameter and a frequency percentage vs. diameter for the sample. From this, the d90 equivalent spherical volume diameter value is derived (90% cumulative undersize value).

20 The compound of structural formula (I) in a free particulate form within the above-mentioned limits, then can be mixed with excipients, diluents, or carriers as necessary to provide, for example, dry powders, aerosols, suspensions, suspension or 25 solid filled capsules, and compressed tablets as oral dosage forms of compound (I).

30 The particle size of free compound (I) in a pharmaceutical composition also can be determined. For example, it is envisioned that the d90 particle size of compound (I) can be determined either in a formulated dosage form or as particles of the free drug, by a microscopic method. First, the composition is separated into its individual

- 15 -

components, or at least compound (I) is separated from the composition. Persons skilled in the art are aware of separation techniques that maintain the particle size of compound (I) during separation of compound (I) from the composition. For example, water-soluble constituents of the composition can be dissolved in water, leaving the highly water insoluble particles of compound (I) without altering the particle size of compound (I) particles.

The undissolved particles then can be examined under a microscope. The crystalline compound (I) can be visually differentiated from amorphous composition ingredients. The particle size of compound (I) is determined by visual inspection and by comparison to standardized particles of a known size. To ensure that the particle size of compound (I) particles is being determined, an infrared microprobe can be used to assay the particles and confirm their identity as compound (I).

Any pharmaceutically acceptable excipients can be used to formulate tablets. The tablets typically contain about 1 to about 20 mg of compound (I). Thus, for example, the particulate compound (I) can be mixed with generally recognized as safe pharmaceutical excipients, including liquid diluents, solid diluents (preferably water-soluble diluents), wetting agents, binders, disintegrants, and lubricants. See, e.g., *Handbook of Pharmaceutical Excipients 2nd Edition*, Amer. Pharm. Assoc. (1994). Preferred solid excipients include lactose, hydroxypropylcellulose, sodium lauryl sulfate, microcrystalline cellulose, talc, colloidal

- 16 -

silicon dioxide, starch, magnesium stearate, stearic acid, and croscarmellose sodium. Liquid excipients include, for example, propylene glycol, glycerin, and ethanol. The pharmaceutical compositions are prepared by standard pharmaceutical manufacturing techniques, as described in *Remington's Pharmaceutical Sciences*, 18th Ed. (1990), Mack Publishing Co., Easton, PA. Such techniques include, for example, wet granulation followed by drying, milling and compression into tablets with or without film coating; dry granulation followed by milling, compression into tablets with or without film coating; dry blending followed by compression into tablets, with or with film coating; molded tablets; sachets; suspensions; wet granulation, dried and filled into gelatin capsules; dry blend filled into gelatin capsules; or suspension filled into gelatin capsules. Generally, solid compositions have identifying marks that are debossed or imprinted on the surface. The total active ingredients in such pharmaceutical compositions comprises from 0.1% to 99.9%, preferably about 1 to 10% by weight of the composition. Preferably, the relative weight percent of excipients is as follows:

- 17 -

	Quantity (% by weight)
Compound (I)	1 to 6
Lactose (diluent)	50 to 75
Hydroxypropylcellulose (binder/diluent)	1 to 5
Croscarmellose Sodium (disintegrant)	3 to 10
Sodium Lauryl Sulfate (wetting agent)	0 to 5
Microcrystalline Cellulose (diluent/disintegrant)	5 to 50
Magnesium Stearate (lubricant)	0.25 to 2.0

15

The specific dose of compound (I) administered according to this invention is, of course, determined by the particular circumstances surrounding the case including, for example, the route of administration, the state of being of the patient, and the pathological condition being treated. A typical daily dose contains a nontoxic dosage level from about 1 to about 20 mg/day of compound (I). Preferred daily doses generally are about 1 to about 20 mg/day, particularly 5 mg, 10 mg, and 20 mg tablets, administered as needed.

The compositions of this invention can be administered by a variety of routes suitable for particulate dosage forms and are preferably administered orally. These compounds preferably are formulated as pharmaceutical compositions prior to administration. The selection of dose is decided by the attending physician.

- 18 -

A compound (I)/hydroxypropylmethylcellulose phthalate coprecipitate was manufactured generally by the method set forth in Butler U.S. Patent No. 5,985,326. After preparation of coprecipitate, the coprecipitate was milled to provide particles having a relatively large particle size and a relatively wide particle size distribution, i.e., $d_{50}=200$ microns. The coprecipitate there was subjected to a controlled dissolution at a pH that ordinarily would not release compound (I) from the polymeric coprecipitate component. Applicants found that the coprecipitate contained a portion of the free drug form of compound (I) not embedded in the coprecipitate polymer. In clinical studies (see Example 2), applicants further discovered that the blood levels of compound (I) within thirty minutes of administration was attributable to the free drug present in the coprecipitate compositions.

These results are surprising in view of Butler U.S. Patent No, 5,985,326 which is directed to a method of preparing a solid dispersion of compound (I) as coprecipitate. The disclosed process and coprecipitate of Butler U.S. Patent No. 5,985,326 is directed to providing a solid dispersion of a poorly water-soluble drug, which has an enhanced bioavailability compared to free particles of the poorly water-soluble drug. Butler U.S. Patent No. 5,985,326 therefore is attempting to avoid the free form of the drug. Butler U.S. Patent No. 5,985,326 generally discloses milling of the coprecipitate, but fails to disclose the size of the coprecipitate particles after milling, and

- 19 -

especially fails to disclose either the presence of the free drug form of compound (I) or, if present, a particle size of the free drug form of compound (I).

Based on these observations, it was concluded that a bimodal delivery of compound (I) could be achieved with a rapid delivery of the free drug followed by a slower delivery of the drug upon the pH sensitive release from the polymeric coprecipitate particles. These observations, in turn, gave rise to the possibility that rapid drug delivery could be effected by compositions incorporating compound (I) entirely in free drug form, provided that suitable stability could be achieved and that the particle size of the drug is controlled in a well-defined range for manufacture of the composition. Accordingly, compound (I) in the pharmaceutical compositions of the present invention preferably is comprised entirely of free drug in particulate form, but alternatively the composition can contain a combination of free drug in particulate form and an embedded drug form to provide a bimodal drug delivery. Preferably, the free drug constitutes greater than 75% free drug (most preferably, greater than 90% free drug) of compound (I) in such compositions.

In one embodiment of the present invention, the free drug form of compound (I), and pharmaceutically-acceptable excipients, diluents, and carriers, are present in a pharmaceutical composition that exhibits a C_{max} (i.e., the maximum observed plasma concentration of compound (I)) of about 180 to about 280 $\mu\text{g/L}$ (micrograms/liter), or an AUC (0-24) (i.e., the area under the plasma

- 20 -

concentration curve from zero to twenty-four hours) of about 2280 to about 3560 $\mu\text{g}\cdot\text{h/L}$ (microgram-hour/liter), measured using a 10 mg dose of the compound. In a preferred embodiment, the composition exhibits 5 a C_{\max} of about 180 to about 280 $\mu\text{g/L}$ and an AUC of about 2280 to about 3650 $\mu\text{g}\cdot\text{h/L}$, measured using a 10 mg dose of the compound. In this embodiment, the 10 composition can be a solid, e.g., a tablet or powder, by using solid diluents, carriers, and/or excipients, or a suspension, e.g., encapsulated in a soft gel, or a solution by using liquid carriers, diluents, and/or carriers.

The C_{\max} and AUC (0-24) were determined by 15 analyzing for compound (I) in plasma using a validated LC/MS/MS method, with a lower limit of quantitation of 0.5 ng/mL. The analytes and an internal standard i.e., the [^{13}C] [$^2\text{H}_3$] isotope of compound (I), were extracted from the plasma by solid phase extraction with 3 mL Empore SD C2 20 cartridges using 150 μL of 90:10 methanol:water. The analytes were separated using high performance liquid chromatography with a Phenomenex Luna phenyl-hexyl (4.6 mm x 100 mm, 5 μ) column with a water: acetonitrile (10:90) mobile phase at 1.0 mL/minute. 25 Detection was performed using a Perkin Elmer Sciex API III Plus tandem mass spectrometer using atmospheric pressure chemical ionization (APCI) in positive ion mode.

It should be understood that C_{\max} and AUC 30 (0-24) in plasma is dose dependent. For example, a composition containing a 20 mg dosage of compound (I) will exhibit a C_{\max} and AUC (0-24) about twice that of a composition containing a 10 mg dosage.

- 21 -

Similarly, a composition containing a 5 mg dosage of compound (I) will exhibit a C_{\max} and AUC (0-24) of about one-half that of a composition containing a 10 mg dosage.

5 Accordingly, the present invention encompasses, for example, compositions containing a 20 mg dosage of compound (I) exhibiting a C_{\max} of about 360 to about 560 $\mu\text{g}/\text{L}$ and/or an AUC (0-24) of about 4560 to about 7120 $\mu\text{g}\cdot\text{h}/\text{L}$; and a composition 10 containing a 5 mg dosage of compound (I) exhibiting a C_{\max} of about 90 to about 140 and/or an AUC (0-24) of about 1140 to about 1780 $\mu\text{g}\cdot\text{h}/\text{L}$. Persons skilled 15 in the art are aware of techniques in which the C_{\max} and AUC (0-24) of compositions containing a dosage of compound (I) different from 10 mg can be compared or standardized to the C_{\max} and AUC (0-24) of a composition containing a 10 mg dose of compound (I).

20 In another embodiment, a composition containing compound (I), either as the free drug alone or as the free drug admixed with a coprecipitate of compound (I), and pharmaceutically-acceptable excipients, diluents, and carriers, exhibits a C_{\max} about 180 to about 280 $\mu\text{g}/\text{L}$ and an AUC (0-24) of about 2280 to about 3650 $\mu\text{g}\cdot\text{h}/\text{L}$. In 25 this embodiment, the composition can be a solid or a suspension.

30 Yet another embodiment of the present invention is a pharmaceutical composition containing a therapeutically-effective amount of particles of compound (I) and pharmaceutically-acceptable carriers, diluents, and excipients, wherein at least 90% of the particles of compound (I) have a particle size of less than about 10 microns, and

- 22 -

bioequivalent compositions thereof. The term "bioequivalent compositions" is defined herein as a composition having a C_{max} of about 180 to about 280 $\mu\text{g}/\text{L}$, and an AUC (0-24) of about 2280 to about 3560 $\mu\text{g}\cdot\text{h}/\text{L}$, measured using a 10 mg dose of particles of compound (I) having a $d90=10$ and a human test subject.

5 C_{max} and AUC (0-24) be determined by methods well-known to person skilled in the art 10 using humans, primates, dogs, rabbits, or rodents (e.g., rats, mice, guinea pigs, and hamsters), for example, as test subjects for bioequivalence. Preferred test animals are humans and dogs.

15 The present invention will be more readily understood upon consideration of the following 20 illustrative examples wherein: Example 1 relates to *in vitro* solubility characteristics of the free drug form of compound (I) of varying particle size; Examples 2 and 3 relate to *in vivo* tests of pharmaceutical compositions incorporating a particulate 25 form according to the invention in comparison to compositions incorporating a coprecipitate and in comparison to compound (I) of a relatively large particle size; and Examples 4 and 5 relate to pharmaceutical compositions employing particulate free drug according to the invention in differing dosage strengths.

EXAMPLE 1

30

In vitro dissolution tests were performed using compound (I) which had been processed by milling from its raw state particulate form ($d90=75-$

- 23 -

200 microns) into particulate preparations having d₉₀ (microns) values as follows: Lot 1, d₉₀=4; Lot 2, d₉₀=22; Lot 3, d₉₀=55; Lot 4, d₉₀=65; Lot 5, d₉₀=73; and Lot 6, d₉₀=116. Alternative milling 5 technologies were employed to develop the various lots. For example, Lot 1 was made using a 12 inch pancake style jet mill fed at a rate of 28-30 kg/hour with sufficient grind pressure to produce the d₉₀=4 material. Lot 2 was prepared in an Alpine VPZ-160 universal mill equipped with pin discs (stud plates) and run at approximately 10,000 rpm. 10

Lots were evaluated *in vitro* by accurately weighing approximately 10 mg of bulk drug into a test tube, adding 1 mL of purified water, and sonicating for up to 2 minutes to ensure the powder was wetted. The drug slurry was subsequently transferred to a dissolution apparatus vessel containing 1000 mL of aqueous 0.5% sodium lauryl sulfate at 15 37°C. The test tube was rinsed with multiple aliquots of warmed dissolution medium and added back into the dissolution vessel. The paddle speed was 50 rpm and samples were taken at 5, 10, 20, and 30 minutes and subsequently analyzed by HPLC. The results are illustrated in Figure 1 and demonstrate 20 improved *in vitro* dissolution occurs with smaller 25 particle sizes of compound (I).

EXAMPLE 2

30 The improvement in bioavailability and reproducibility of pharmaceutical compositions made available by the present invention is demonstrated *in vivo* in humans. The following Table 1 demon-

- 24 -

strates the pharmaceutical compositions prepared as in Examples 4 and 5 with particulate free drug having a d₉₀ of 8.4 microns compared to composition incorporating the coprecipitate of compound (I) with hydroxypropylmethylcellulose phthalate (coprecipitate). In each instance, the tableted composition was designed to deliver a 10 mg dose of compound (I).

10

15

Table 1 In vivo evaluation		
Pharmaceutical Composition	No. of Patients	T _{max} (hrs)
Free Drug of Compound (I)	18	2.0
Coprecipitate of Compound (I)	18	3.5

20 The composition incorporating a particu-
late free drug form having a d₉₀ of 8.4 demonstrated
significantly improved T_{max} over a composition con-
taining the coprecipitate (T_{max} is a measure of the
time to achieve peak blood levels of a drug, and is
indicative of improved onset of action). The
25 particulate free drug formulation correspondingly
provided a more rapid rate of absorption of compound
(I) into plasma, providing a geometric mean plasma
level at 30 minutes of 51 ng/ml (nanograms per
milliliter) as compared to 29 ng/ml for the
30 coprecipitate formulation.

- 25 -

EXAMPLE 3

5 A study was conducted to determine the bioequivalence of tablets containing compound (I) in different particle sizes. The tablets contained compound (I) in a particle size of $d_{90}=8.4\mu$ (micron), $d_{90}=20\mu$, or $d_{90}=52\mu$.

10 The study was an open-label, randomized, three-period crossover study conducted on twenty-four (24) healthy male subjects aged 18 to 65 years old, divided into two groups of twelve. A single 10 mg oral dose was administered with 180 mL of water in each of three treatment periods, and the 15 pharmacokinetics of tablets containing compound (I) in different particle sizes were compared.

20 After dosing, the subjects underwent pharmacokinetic blood sampling. There was an interval of at least 10 days between dosing in each treatment period to eliminate any residual compound (I) from the previous treatment period. The post-study assessment was conducted between 7 and 14 days after the final dosing.

25 Compound (I) was absorbed relatively quickly following oral dosing from the $d_{90}=52$, 30, and 8.4μ particle size formulations. However, the rate and extent of absorption of compound (I) increased with decreasing particle size. A comparison of C_{max} and AUC (0-24) data showed that the difference in absorption between particle size formulations was most apparent over the first 24 hours after dosing. As used herein, C_{max} is defined as the maximum observed plasma concentration of compound 30

- 26 -

(I), and AUC (0-24) is defined as the area under the plasma concentration time curve from zero to twenty-four hours. Both C_{max} and AUC (0-24) are well-known and understood variables to persons skilled in the art.

With respect to C_{max} , the $d90=52\mu$ and $d90=20\mu$ formulations were not bioequivalent to the $d90=8.4\mu$ formulation because the 90% confidence interval (CI) was outside of the 0.8 to 1.25 equivalence limits. In particular, C_{max} was 36% and 23% lower for the 52μ and 20μ formulations, respectively, compared to the 8.4μ formulation. The 52μ formulation also was not equivalent to the 8.4μ formulation with respect to AUC (0-24), which was 23% lower than the 8.4μ formulation. The 20μ and 8.4μ formulations were bioequivalent with respect to AUC (0-24). The 8.4μ , 20μ , and 52μ formulations were bioequivalent with respect to AUC, i.e., the area under the plasma concentration time curve from time zero to infinity.

The study showed that the rate of absorption of compound (I), based on C_{max} and t_{max} (i.e., time to attain maximum observed drug-plasma concentration), was slower for the 52μ formulations in relation to the 8.4μ formulation. As stated above, C_{max} was not equivalent for the 52μ and 20μ formulations compared to the 8.4μ formulation. Median t_{max} occurred one hour later for the 52μ formulation, but was similar to the 20μ and 8.4μ formulations.

The following table summarizes various pharmacokinetic parameters of compound (I) following

- 27 -

oral administration of a single 10 mg dose of the d90 52 μ , 20 μ , and 8.4 μ particle size formulations.

	d90=52 μ	d90=20 μ	d90=8.4 μ	
5	C _{max} (μ g/L)	142	189	224
	t _{max} (h) ¹⁾	3.00	2.00	2.00
	AUC (0-24) ²⁾	2201	2667	2849

10 ¹⁾ median data.

²⁾ in micrograms·hour/liter.

15 This study showed that reducing the par-
ticle size of compound (I) in accordance with the
present invention has an impact on the *in vivo* rate
of absorption of compound (I) from a solid dosage
form, and, hence, on the bioavailability of compound
(I). For example, from the statistical analysis,
20 t_{max} for the 52 μ formulation occurred significantly
(i.e., 1 hour) later than for the 8.4 μ formulation.
There was no significant difference in t_{max} between
the 20 μ and 8.4 μ formulations. Accordingly, onset
25 of a therapeutic benefit attributed to compound (I)
after administration is significantly faster for the
8.4 μ and 20 μ formulations compared to the 52 μ form-
ulation.

30 In addition to dissolution and *in vivo*
absorption, another important aspect of the physical
properties of particulate β -carboline preparations
according to the present invention is the impact on
the various unit operations of the drug product
manufacturing process. While the particle size
35 specification ensures consistent delivery of the

- 28 -

drug molecule to the sites of absorption in the gastrointestinal tract, it also imparts better control during the tablet manufacturing process.

5 The following formulation examples are illustrative only and are not intended to limit the scope of the present invention.

EXAMPLE 4

10 The following formula was used to prepare the finished dosage form of a tablet providing 10 mg of compound (I).

Ingredient	Quantity (mg)
Granulation	
Compound (I) (Lot 1, d90 of 4)	10.00
Lactose Monohydrate	153.80
Lactose Monohydrate (Spray Dried)	25.00
Hydroxypropylcellulose (EF Extra Fine)	4.00
20 Croscarmellose Sodium	9.00
Hydroxypropylcellulose (EF)	1.75
Sodium Lauryl Sulfate	0.70
Outside Powders	
Microcrystalline Cellulose (Granular-102)	37.50
25 Croscarmellose Sodium	7.00
Magnesium Stearate (Vegetable)	1.25
Total	250 mg

30 Purified Water, USP was used in the manufacture of the tablets. The water was removed during processing and minimal levels remained in the finished tablets.

- 29 -

Tablets are manufactured using a wet granulation process. A step-by-step description of the process follows. Compound (I) and excipients to be granulated are security sieved. Compound (I) is dry blended with lactose monohydrate (spray dried), hydroxypropylcellulose, croscarmellulose sodium, and lactose monohydrate. The resulting powder blend was granulated with an aqueous solution of hydroxypropylcellulose and sodium lauryl sulfate using a Powrex or other suitable high shear granulation. Additional water can be added to reach the desired endpoint. A mill can be used to delump the wet granulation and facilitate drying. The wet granulation was dried using either a fluid bed dryer or a drying oven. After drying, the material can be sized to eliminate any large agglomerates. Microcrystalline cellulose, croscarmellose sodium, and magnesium stearate were security sieved and added to the dry sized granules. These excipients and the dry granulation were mixed until uniform using a tumble bin, ribbon mixer, or other suitable mixing equipment. The mixing process can be separated into two phases. The microcrystalline cellulose, croscarmellose sodium, and the dried granulation were added to the mixer and blended during the first phase, followed by the addition of the magnesium stearate to this granulation and a second mixing phase.

The mixed granulation then was compressed into tablets using a rotary compression machine. The core tablets were film coated with an aqueous suspension of the appropriate color mixture in a coating pan (e.g., Accela Cota). The coated tablets

- 30 -

can be lightly dusted with talc to improve tablet handling characteristics.

5 The tablets are filled into plastic containers (30 tablets/container) and accompanied by package insert describing the safety and efficacy of the formulation.

EXAMPLE 5

10 By analogous procedures, the following formula was used to prepare the finished dosage form of a tablet providing 5.0 mg and 20 mg of compound (I).

	Ingredient	Quantity (mg)	Quantity (mg)
Granulation			
	Compound (I) (Lot 1, d90 of 4)	5.00	20.00
	Lactose Monohydrate	109.66	210.19
	Lactose Monohydrate (Spray Dried)	17.50	35.00
20	Hydroxypropylcellulose	2.80	5.60
	Croscarmellose Sodium	6.30	12.60
	Hydroxypropylcellulose (EF)	1.22	2.45
	Sodium Lauryl Sulfate	0.49	0.98
Outside Powders			
25	Microcrystalline Cellulose (Granular-102)	26.25	52.50
	Croscarmellose Sodium	4.90	9.80
	Magnesium Stearate (Vegetable)	0.88	0.88
30	Total	175 mg	350 mg


The principles, preferred embodiments, and modes of operation of the present invention have

- 31 -

been described in the foregoing specification. The invention that is intended to be protected herein, however, is not construed to be limited to the particular forms disclosed, because they are to be regarded as illustrative rather than restrictive. Variations and changes may be made by those skilled in the art without departing from the spirit of the invention.

WHAT IS CLAIMED IS:

1. A free drug particulate form of a compound having a formula



and pharmaceutically acceptable salts and solvates thereof, comprising particles of the compound wherein at least 90% of the particles have a particle size of less than about 40 microns.

2. The free drug particulate form of claim 1 wherein at least 90% of the particles have a particle size of less than about 25 microns.

3. The free drug particulate form of claim 1 wherein at least 90% of the particles have a particle size of less than about 15 microns.

4. The free drug particulate form of claim 1 wherein at least 90% of the particles have a particle size of less than about 10 microns.

- 33 -

5. A pharmaceutical composition comprising the free drug particulate form of claim 1 and one or more pharmaceutically-acceptable carriers, diluents, or excipients.

6. The pharmaceutical composition of claim 5 wherein the free drug is entirely in particulate form.


7. A method of treating sexual dysfunction in a patient in need thereof, which comprises administering to the patient a therapeutically effective amount of a composition comprising the free drug particulate form of claim 1 and one or more pharmaceutically-acceptable carriers, diluents, or excipients.

8. The method of claim 7 wherein the sexual dysfunction is male erectile dysfunction.

9. The method of claim 7 wherein the sexual dysfunction is female sexual arousal disorder.

- 34 -

10. A pharmaceutical composition comprising:
(a) a free drug form of a compound having the formula



and pharmaceutically-acceptable salts and solvates thereof, and


(b) one or more pharmaceutically-acceptable carriers, diluents, or excipients, wherein the composition exhibits a C_{max} of about 180 to about 280 micrograms/liter or an AUC (0-24) of about 2280 to about 3560 microgram · hour/liter, measured using a 10 milligram dose of the compound.

11. The composition of claim 10 wherein the composition exhibits a C_{max} of about 180 to about 280 micrograms/liter and an AUC (0-24) of about 2280 to about 3560 microgram · hour/liter.

- 35 -

12. A pharmaceutical composition comprising:

(a) a compound having the formula




and pharmaceutically-acceptable salts and solvates thereof, and

(b) one or more pharmaceutically-acceptable carriers, diluents, or excipients, wherein the composition exhibits a C_{max} of about 180 to about 280 micrograms/liter and an AUC (0-24) of about 2280 to about 3560 micrograms · hour/liter, measured using a 10 milligram dose of the compound.

- 36 -

13. A pharmaceutical composition comprising:

(a) a free drug form of a compound having the formula



and pharmaceutically acceptable salts and solvates thereof, wherein at least 90% of the particles have a particle size of less than about 10 microns, and

(b) one or more pharmaceutically-acceptable carriers, diluents, or excipients, and bioequivalent compositions thereof.

14. A method of manufacturing the free drug particulate form of claim 1 comprising:


(a) providing a solid, free form of the compound, and

(b) comminuting the solid free form of the compound to provide particles of the compound wherein at least 90% of the particles have a particle size of less than about 40 microns.

- 37 -

15. The method of claim 14 further comprising the step of admixing the particles of step (b) with one or more pharmaceutically-acceptable carriers, diluents, or excipients.


16. A pharmaceutical composition prepared by admixing particles of a compound having a formula



or a pharmaceutically-acceptable salt or solvate thereof, with one or more pharmaceutically-acceptable carrier, diluent, or excipient, wherein the particles of the compound have a d90=40 or less.

- 38 -


17. Use of particles of a free form of a compound having a formula



wherein at least 90% of the particles have a particle size of less than about 40 microns, for the manufacture of a medicament for the treatment of male erectile dysfunction or female sexual arousal disorder.

- 39 -

18. Use of particles of a free form of a compound having a formula



which exhibits a C_{max} of about 180 to about 280 micrograms/liter and an AUC (0-24) of about 2280 to about 3560 micrograms·hour/liter, measured using an 10 milligram dose of the compound, for the manufacture of a medicament for the treatment of male erectile dysfunction or female arousal disorder.

19. The invention as herein above described.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
8 February 2001 (08.02.2001)

PCT

(10) International Publication Number
WO 01/08688 A3

(51) International Patent Classification⁷: A61K 31/4985, 9/14, A61P 15/10

(21) International Application Number: PCT/US00/20981

(22) International Filing Date: 1 August 2000 (01.08.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data: 60/147,048 3 August 1999 (03.08.1999) US

(71) Applicant (for all designated States except US): **LILLY ICOS LLC** [US/US]; 1209 Orange Street, Wilmington, DE 19801 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **ANDERSON, Neil, R.** [US/US]; 906 Northwestern Avenue, West Lafayette, IN 47906 (US). **HARTAUER, Kerry, J.** [US/US]; 14907 Pacer Court, Carmel, IN 46032 (US). **KRAL, Martha, A.** [US/US]; 6125 Haverford Avenue, Indianapolis, IN 46220 (US). **STEPHENSON, Gregory, A.** [US/US]; 10227 Bent Tree Lane, Fishers, IN 46032 (US).

(74) Agent: **NAPOLI, James, J.**; Marshall, O'Toole, Gerstein, Murray & Borun, 6300 Sears Tower, 233 S. Wacker Drive, Chicago, IL 60606 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).



Published:

— with international search report

(88) Date of publication of the international search report: 16 August 2001

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.


(54) Title: BETA-CARBOLINE DRUG PRODUCTS



WO 01/08688 A3

(57) Abstract: A compound of structural formula (I), and pharmaceutically acceptable salts and solvates thereof, wherein the compound is in free drug particulate form, is disclosed.

Figure 1
Comparison of Bulk Drug Substance Dissolution Profiles



DECLARATION FOR PATENT APPLICATION AND POWER OF ATTORNEY

As a below named inventor, I hereby declare that my residence, post office address and citizenship are as stated below next to my name; I believe that I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled "**β-CARBOLINE DRUG PRODUCTS**," the specification of which (check one): is attached hereto; was filed on _____ as Application Serial No. _____ and was amended on _____ (if applicable); was filed as PCT International Application No. PCT/US00/20981 on August 1, 2000, and was amended under Article 19 on _____ (if applicable). I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment(s) referred to above. I acknowledge the duty to disclose to the Patent and Trademark Office all information known to me to be material to patentability as defined in 37 C.F.R. §1.56.

I hereby claim foreign priority benefits under 35 U.S.C. §119 of any foreign application(s) for patent or inventor's certificate or of any PCT international application(s) designating at least one country other than the United States of America listed below and have also identified below any foreign application(s) for patent or inventor's certificate or any PCT international application(s) designating at least one country other than the United States of America filed by me on the same subject matter having a filing date before that of the application(s) of which priority is claimed:

			Priority Claimed
PCT/US00/20981 (Application Serial Number)	PCT (Country)	01/08/00 (Day/Month/Year Filed)	<input checked="" type="checkbox"/> <input type="checkbox"/> Yes No
_____ (Application Serial Number)	_____ (Country)	_____ (Day/Month/Year Filed)	<input type="checkbox"/> <input checked="" type="checkbox"/> Yes No

I hereby claim the benefit under 35 U.S.C. §119(e) of any United States provisional application(s) listed below:

60/147,048 (Application Serial Number)	01/08/99 (Day/Month/Year Filed)
_____ (Application Serial Number)	_____ (Day/Month/Year Filed)

I hereby claim the benefit under 35 U.S.C. §120 of any United States application(s) or PCT international application(s) designating the United States of America listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior application(s) in the manner provided by the first paragraph of 35 U.S.C. §112, I acknowledge the duty to disclose to the Office all information known to me to be material to patentability as defined in 37 C.F.R. §1.56 which occurred between the filing date of the prior application(s) and the national or PCT international filing date of this application:

(Application Serial Number)	(Day/Month/Year Filed)	(Status-Patented, Pending or Abandoned)
_____ (Application Serial Number)	_____ (Day/Month/Year Filed)	_____ (Status-Patented, Pending or Abandoned)

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under 18 U.S.C. §1001 and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

POWER OF ATTORNEY: I hereby appoint as my attorneys, with full powers of substitution and revocation to prosecute this application and transact all business in the Patent and Trademark Office connected therewith:

John B. Lungmus (18,566)
 Allen H. Gerstein (22,218)
 Nate F. Scarpelli (22,320)
 Michael F. Borun (25,447)
 Carl E. Moore, Jr. (26,487)
 Richard H. Anderson (26,526)

Patrick D. Ertel (26,877)
 Richard B. Hoffman (26,910)
 James P. Zeller (28,491)
 Kevin D. Hogg (31,839)
 Jeffrey S. Sharp (31,879)
 Martin J. Hirsch (32,237)

James J. Napoli (32,361)
 Richard M. La Barge (32,254)
 Robert M. Gerstein (34,824)
 Anthony G. Sitko (36,278)
 James A. Flight (37,622)

Roger A. Heppermann (37,641)
 David A. Gass (38,153)
 Gregory C. Mayer (38,238)
 Michael R. Weiner (38,359)
 William K. Merkel (40,725)

Send correspondence to: James J. Napoli

FIRM NAME	PHONE NO.	STREET	CITY & STATE	ZIP CODE
Marshall, Gerstein & Borun	312-474-6300	6300 Sears Tower 233 South Wacker Drive	Chicago, Illinois	60606-6357

Full Name of First or Sole Inventor <i>100</i> Neil R. Anderson	Citizenship United States of America
Residence Address - Street 906 Northwestern Avenue	Post Office Address - Street 906 Northwestern Avenue
City (Zip) West Lafayette (47906) <i>IN</i>	City (Zip) West Lafayette (47906)
State or Country Indiana	State or Country Indiana
Date <input checked="" type="checkbox"/> 1-3-2002	Signature <i>Neil Anderson</i>

Second Joint Inventor, if any <i>200</i> Kerry J. Hartauer	Citizenship United States of America
Residence Address - Street 14907 Pacer Court	Post Office Address - Street 14907 Pacer Court
City (Zip) Carmel (46032) <i>IN</i>	City (Zip) Carmel (46032)
State or Country Indiana	State or Country Indiana
Date <input checked="" type="checkbox"/> 1-16-02	Signature <i>Kerry J. Hartauer</i>

Third Joint Inventor, if any <i>300</i> Martha A. Kral	Citizenship United States of America
Residence Address - Street 6125 Haverford Avenue	Post Office Address - Street 6125 Haverford Avenue
City (Zip) Indianapolis (46220) <i>IN</i>	City (Zip) Indianapolis (46220)
State or Country Indiana	State or Country Indiana
Date <input checked="" type="checkbox"/> 1-16-02	Signature <i>Martha A. Kral</i>

Fourth Joint Inventor, if any <i>400</i> Gregory A. Stephenson	Citizenship United States of America
Residence Address - Street 10227 Bent Tree Lane	Post Office Address - Street 10227 Bent Tree Lane
City (Zip) Fishers (46032) <i>IN</i>	City (Zip) 800.4.1 Fishers (46032)
State or Country Indiana	State or Country Indiana
Date <input checked="" type="checkbox"/> 1-16-02	Signature <i>Gregory A. Stephenson</i>